



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2020

---

## **Clinical Imaging of the Heterogeneous Group of Triple-negative Breast Cancer**

Müller, Martin ; Güth, Uwe ; Varga, Zsuzsanna ; Reeve, Kelly ; Bjelic-Radisic, Vesna ; Fleisch, Markus ; Tausch, Christoph J ; Elfgen, Constanze

**Abstract:** BACKGROUND/AIM Triple-negative breast cancer (TNBC) can be divided into subtypes of basal-like (BL), mesenchymal-like (ML), luminal androgen receptor (LAR), and immunomodulatory (IM). The aim of our study was to assess whether there are distinct radiologic features within the different TNBC subtypes and whether this has potential clinical impact. **PATIENTS AND METHODS** Imaging pictures of 135 patients with TNBC were re-evaluated. TNBC subtyping was performed on asservated tumor tissue using a panel of antibodies. **RESULTS** Mammographic margins of LAR-TNBC were more often spiculated (24.3% versus 0-4.1%). BL-TNBC presented more frequent a mass without calcification in mammogram than other subtypes (71.4% versus 48.6-57.9%). In ultrasound, ML and LAR were described more often with smooth borders. **CONCLUSION** The histopathological subtype of TNBC influences its presentation in ultrasound and mammogram. This can reflect a different growth pattern of the subtypes and may have an impact on the early diagnosis of TNBC.

DOI: <https://doi.org/10.21873/anticancerres.14171>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-186896>

Journal Article

Published Version

Originally published at:

Müller, Martin; Güth, Uwe; Varga, Zsuzsanna; Reeve, Kelly; Bjelic-Radisic, Vesna; Fleisch, Markus; Tausch, Christoph J; Elfgen, Constanze (2020). Clinical Imaging of the Heterogeneous Group of Triple-negative Breast Cancer. *Anticancer Research*, 40(4):2125-2131.

DOI: <https://doi.org/10.21873/anticancerres.14171>

## Clinical Imaging of the Heterogeneous Group of Triple-negative Breast Cancer

MARTIN MÜLLER<sup>1</sup>, UWE GÜTH<sup>1</sup>, ZSUZSANNA VARGA<sup>2</sup>, KELLY REEVE<sup>3</sup>, VESNA BJELIC-RADISIC<sup>4,5</sup>,  
MARKUS FLEISCH<sup>4,5</sup>, CHRISTOPH J. TAUSCH<sup>1</sup> and CONSTANZE ELFGEN<sup>1,5</sup>

<sup>1</sup>Breast Center Zurich, Zurich, Switzerland;

<sup>2</sup>Institute of Pathology and Molecular Pathology, University Hospital of Zurich, Zurich, Switzerland;

<sup>3</sup>Epidemiology, Biostatistics and Prevention Institute, Biostatistics Department,  
University of Zurich, Zurich, Switzerland;

<sup>4</sup>Department for Gynecology and Obstetrics, Helios Hospital Wuppertal, Wuppertal, Germany;

<sup>5</sup>University of Witten-Herdecke, Witten-Herdecke, Germany

**Abstract.** *Background/Aim:* Triple-negative breast cancer (TNBC) can be divided into subtypes of basal-like (BL), mesenchymal-like (ML), luminal androgen receptor (LAR), and immunomodulatory (IM). The aim of our study was to assess whether there are distinct radiologic features within the different TNBC subtypes and whether this has potential clinical impact. *Patients and Methods:* Imaging pictures of 135 patients with TNBC were re-evaluated. TNBC subtyping was performed on asservated tumor tissue using a panel of antibodies. *Results:* Mammographic margins of LAR-TNBC were more often spiculated (24.3% versus 0-4.1%). BL-TNBC presented more frequent a mass without calcification in mammogram than other subtypes (71.4% versus 48.6-57.9%). In ultrasound, ML and LAR were described more often with smooth borders. *Conclusion:* The histopathological subtype of TNBC influences its presentation in ultrasound and mammogram. This can reflect a different growth pattern of the subtypes and may have an impact on the early diagnosis of TNBC.

Molecular typing led to a vast improvement in the diagnosis and treatment of breast cancer (BC) decades ago. Targeted therapy was developed for cancers with hormonal receptor positivity and for HER2 positivity, which significantly improved their prognosis. Lacking the three markers of estrogen receptor (ER), progesterone receptor (PR), and HER2 receptor, triple negative breast cancer (TNBC) remains the most feared diagnosis with poor outcomes (1). TNBC is

associated with younger age and a higher tumor stage at diagnosis, as well as BRCA 1 germline mutation (2). Today, TNBC is regarded as a heterogeneous disease. Subtyping provides further information about the molecular pathways, chemotherapy response, and the potential for development of targeted therapies (3). The TNBC subtypes have been classified into basal-like (BL), mesenchymal-like (ML), and luminal androgen receptor (LAR); lymphocytic infiltration defines a further subtype of immunomodulatory (IM) TNBC (3-5).

Early detection is crucial for the curative treatment of BC, especially for aggressive tumors. Mammography is considered to be the gold standard for screening of BC. In cases of dense breast tissue, family history of BC, and in women with suspicious findings in mammogram or clinical examination, additional ultrasound is indicated. For the objective assessment of breast lesions, the Breast Imaging Reporting and Data System (BI-RADS) has been established (6). Several features of mammographic and sonographic breast findings are evaluated algorithmically to minimize radiologist-dependent subjective ratings. TNBC is known to present more often benign characteristics in clinical imaging such as lobulated, oval-shaped, and circumscribed margins, than other types of BC (7). Benign classification and misinterpretation can lead to diagnostic delay and negative impact on the outcomes of BC (8). Despite TNBC subtypes having different therapy responses and outcomes, very little is known about their characteristics in clinical imaging. In this study we assessed specific features of TNBC subtypes in mammogram and ultrasound with the aim of improving the understanding of this heterogeneous disease.

*Correspondence to:* Constanze Elfgen, MD, Breast Center Zurich, Seefeldstrasse 214, 8008 Zurich, Switzerland. Tel: +41 445338100, e-mail: c.elfgen@brust-zentrum.ch

**Key Words:** Triple negative breast cancer, TNBC subtypes, ultrasound, clinical imaging, breast cancer.

### Patients and Methods

Patients with TNBC who were diagnosed in the Breast-Center Zurich between 2002 and 2016 were included in the study. Patient- and disease-related data as well as outcome were evaluated retrospectively.

Table I. Patient and tumor characteristics.

| n                                       | Overall<br>135 | BL<br>49    | IM<br>19    | ML<br>30    | LAR<br>37   | p-Value |
|---|----------------|-------------|-------------|-------------|-------------|---------|
| Age [y; mean (sd)]                      | 57.7 (13.7)    | 55.3 (14.4) | 54.7 (14.4) | 57.0 (12.4) | 62.9 (12.3) | 0.07    |
| Menopausal status (%)                   |                |             |             |             |             | 0.02    |
| Premenopausal                           | 44 (32.6)      | 20 (40.8)   | 9 (47.4)    | 10 (33.3)   | 5 (13.5)    |         |
| Postmenopausal                          | 91 (67.4)      | 29 (59.2)   | 10 (52.6)   | 20 (66.7)   | 32 (86.5)   |         |
| Reason for first consultation (%)       |                |             |             |             |             | 0.39    |
| Self-detected palpable mass             | 80 (59.3)      | 35 (71.4)   | 12 (63.2)   | 18 (60.0)   | 15 (40.5)   |         |
| Screening                               | 41 (30.4)      | 10 (20.4)   | 6 (31.6)    | 11 (36.7)   | 14 (37.8)   |         |
| Symptoms (discharge/pain)               | 5 (3.7)        | 1 (2.0)     | 0           | 1 (3.3)     | 3 (8.1)     |         |
| other                                   | 9 (6.7)        | 3 (6.1)     | 1 (5.3)     | 0           | 5 (13.5)    |         |
| Family history of breast cancer (%)     |                |             |             |             |             | 0.90    |
| Yes                                     | 24 (17.8)      | 9 (18.4)    | 4 (21.1)    | 6 (20.0)    | 5 (13.5)    |         |
| No                                      | 110 (81.5)     | 40 (81.6)   | 15 (78.9)   | 24 (80.0)   | 31 (83.8)   |         |
| Unknown                                 | 1 (0.7)        | 0           | 0           | 0           | 1 (2.7)     |         |
| Histopathologic tumor size (%)          |                |             |             |             |             | 0.98    |
| ypT0                                    | 5 (3.7)        | 2 (4.1)     | 1 (5.3)     | 1 (3.3)     | 1 (2.7)     |         |
| ypT1                                    | 19 (14.1)      | 6 (12.2)    | 5 (26.3)    | 4 (13.3)    | 4 (10.8)    |         |
| ypT2-4                                  | 8 (5.9)        | 2 (4.1)     | 1 (5.3)     | 3 (10.0)    | 2 (5.4)     |         |
| pT1                                     | 45 (33.3)      | 16 (32.7)   | 8 (42.1)    | 10 (33.3)   | 11 (29.7)   |         |
| pT2-4                                   | 58 (43.0)      | 23 (46.9)   | 4 (21.1)    | 12 (40.0)   | 19 (51.4)   |         |
| Histopathologic axillary lymph node (%) |                |             |             |             |             | 0.44    |
| ypN0                                    | 13 (9.6)       | 5 (10.2)    | 4 (21.1)    | 1 (3.3)     | 3 (8.1)     |         |
| ypN1-3                                  | 14 (10.4)      | 3 (6.1)     | 3 (15.8)    | 4 (13.3)    | 4 (10.8)    |         |
| pN0                                     | 77 (57.0)      | 30 (61.2)   | 10 (52.6)   | 19 (63.3)   | 18 (48.6)   |         |
| pN1-3                                   | 31 (23.0)      | 11 (22.4)   | 2 (10.5)    | 6 (20.0)    | 12 (32.4)   |         |
| Tumor grading (%)                       |                |             |             |             |             | 0.05    |
| Well differentiated (G1-2)              | 21 (15.6)      | 4 (8.2)     | 3 (15.8)    | 3 (10.0)    | 11 (29.7)   |         |
| Poor differentiated (G3)                | 114 (84.4)     | 45 (91.8)   | 16 (84.2)   | 27 (90.0)   | 26 (70.3)   |         |

At patient's first presentation, initial evaluation had been performed and documented by trained breast radiologists in the Breast-Center Zurich. Standardized mammogram had been performed in two imaging planes (mediolateral oblique and craniocaudal). Ultrasound with 5-12 MHz transducers had been performed according to the BI-RADS guidelines. For classification reassessment, mammogram and ultrasound pictures were retrospectively reviewed by two experienced breast radiologists who were blinded for the patient's history and histopathological subtyping of the tumor. Radiological tumor characteristics such as margin, shape, orientation, echo pattern, and posterior features were categorized according to the BI-RADS classification. To analyse potential factors for misdiagnosis, the initially given BI-RADS classification (benign/probably benign *versus* probably malignant) was not changed but correlated to imaging features.

At initial diagnosis, surgical and biopsy specimens had been processed as formalin-fixed, paraffin-embedded tumor tissues according to the standardized protocol of the Department of Pathology and Molecular Pathology, University Hospital Zurich, Switzerland. TNBC had been defined as expressing less than 1% of ER and PR by immunohistochemistry and being negative for HER2 expression. A tissue micro array (TMA) was constructed from the asservated surgical specimen according to Kündig *et al.* as described previously (9, 10). TNBC subtyping into Basel-like (BL), Immunomodulatory (IM), Mesenchymal-like (ML), and Luminal androgen receptor (LAR) was performed using a panel of antibodies according to Lehmann *et al.* and Turner *et al.* (3, 5).

For the descriptive analysis, mean (and standard deviation) or median were used for continuous variables and number and percentage were used for categorical variables. To enhance interpretation, stacked bar plots were used to visualize some categorical variables. Chi square and Fisher's exact test were used to test for differences in proportions across subtypes. Overall survival and disease-free survival were visualized with Kaplan-Meier plots of survival probability. All analyses were performed in the R programming language.

The study was approved by the local ethics committee of Zurich, Switzerland (BASEC-No. 2017-00219), according to the national and international ethics guidelines. Informed consent was obtained from all individual participants included in the study.

## Results

We collected data from a total of 166 patients with TNBC. For the present study, 31 patients had to be excluded due to the low quality of digital stored images that did not allow a re-evaluation. The average age of the patients was 58 years (sd=14) and was similar across the subtypes (Table I). A higher percentage of postmenopausal women was found in the LAR subtype. Most patients reported a self-detected palpable tumor as the reason for first consultation, followed

Table II. Imaging characteristics in ultrasound and mammogram of distinct TNBC subtypes according to BI-RADS classification.

| n   | Overall<br>135 | BL<br>49  | IM<br>19  | ML<br>30  | LAR<br>37 | p-Value |
|---|----------------|-----------|-----------|-----------|-----------|---------|
| Sonographic shape (%)                                 |                |           |           |           |           | 0.26    |
| Round/oval  | 6 (4.4)        | 1 (2.0)   | 2 (10.0)  | 1 (3.3)   | 2 (5.4)   |         |
| Irregular   | 125 (92.6)     | 48 (98.0) | 17 (10.5) | 27 (90.0) | 33 (89.2) |         |
| No detected tumor                                     | 4 (3.0)        | 0         | 0         | 2 (6.7)   | 2 (5.4)   |         |
| Sonographic margin (%)                                |                |           |           |           |           | 0.11    |
| Circumscribed   | 11 (8.1)       | 3 (6.1)   | 3 (15.8)  | 1 (3.3)   | 4 (10.8)  |         |
| Spiculated  | 71 (52.6)      | 30 (61.2) | 13 (68.4) | 14 (46.7) | 14 (37.6) |         |
| Lobulated   | 49 (36.3)      | 16 (32.7) | 3 (15.8)  | 13 (43.3) | 17 (45.9) |         |
| No detected tumor                                     | 4 (3.0)        | 0         | 0         | 3 (6.7)   | 2 (5.4)   |         |
| Sonographic pattern (%)                               |                |           |           |           |           | 0.44    |
| Complex   | 21 (15.6)      | 9 (18.4)  | 5 (26.3)  | 2 (6.7)   | 5 (13.5)  |         |
| Hypoechoic  | 109 (80.7)     | 39 (79.6) | 14 (73.7) | 26 (86.7) | 30 (81.1) |         |
| Isoechoic   | 1 (0.7)        | 1 (2.0)   | 0         | 0         | 0         |         |
| No detected tumor                                     | 4 (3.0)        | 0         | 0         | 2 (6.7)   | 2 (5.4)   |         |
| Sonographic posterior pattern (%)                     |                |           |           |           |           | 0.031   |
| Posterior enhancement                                 | 67 (49.6)      | 28 (57.1) | 11 (57.9) | 17 (56.1) | 11 (29.7) |         |
| Posterior shadowing                                   | 34 (25.2)      | 13 (26.5) | 1 (5.3)   | 7 (23.3)  | 13 (35.1) |         |
| No change   | 30 (22.2)      | 8 (16.3)  | 7 (36.8)  | 4 (13.3)  | 11 (29.7) |         |
| No detected tumor                                     | 4 (3.0)        | 0         | 0         | 2 (6.7)   | 2 (5.4)   |         |
| Sonographic orientation                               |                |           |           |           |           | 0.98    |
| Horizontal  | 90 (66.7)      | 35 (71.4) | 13 (68.4) | 18 (60.0) | 24 (64.9) |         |
| Vertical  | 30 (22.2)      | 10 (20.4) | 4 (21.1)  | 7 (23.3)  | 9 (24.3)  |         |
| Round   | 11 (8.1)       | 4 (8.2)   | 2 (10.5)  | 3 (10.0)  | 2 (5.4)   |         |
| No detected tumor                                     | 4 (3.0)        | 0         | 0         | 2 (6.7)   | 2 (5.4)   |         |
| Mammographic shape                                    |                |           |           |           |           | 0.007   |
| Focal asymmetry                                       | 12 (8.9)       | 4 (8.2)   | 1 (5.3)   | 4 (13.3)  | 3 (8.1)   |         |
| Mass  | 79 (58.5)      | 35 (71.4) | 11 (57.9) | 15 (50.0) | 18 (48.6) |         |
| Mass with calcification                               | 18 (13.3)      | 3 (6.1)   | 1 (5.3)   | 3 (10.0)  | 11 (29.7) |         |
| Calcification only                                    | 7 (5.2)        | 0         | 0         | 4 (13.3)  | 3 (8.1)   |         |
| Distortion  | 0              | 0         | 0         | 0         | 0         |         |
| No detected tumor                                     | 19 (14.1)      | 7 (14.3)  | 6 (31.6)  | 4 (13.3)  | 2 (5.4)   |         |
| Mammographic margins                                  |                |           |           |           |           | 0.077   |
| Circumscribed   | 51 (37.8)      | 21 (42.9) | 7 (36.8)  | 11 (36.7) | 12 (32.4) |         |
| Spiculated  | 12 (8.9)       | 2 (4.1)   | 0         | 1 (3.3)   | 9 (24.3)  |         |
| Indistinct  | 46 (34.1)      | 19 (38.8) | 6 (31.6)  | 10 (33.3) | 11 (29.7) |         |
| Not applicable (no detected tumor/calcification only) | 26 (19.3)      | 7 (14.3)  | 6 (31.6)  | 8 (26.7)  | 5 (13.5)  |         |

by findings from a referring physician and findings during screening at the Breast Center (Table I). Tumor stage did not vary across subtypes. LAR tumors appeared slightly more often well differentiated than other subtypes (29.7%; n=11).

Breast density did not significantly differ within the subtypes. No tumor was seen upon ultrasound in four patients (3.0%), while mammography failed to detect 19 cases of TNBC (14.1%). Benign sonographic classification was given in 19 cases (14.5%). Most of these showed a horizontal orientation, hypoechoic pattern, and posterior enhancement (Table II, Figures 1 and 2). Overall, 66.7% of tumors displayed a horizontal orientation, 22.2% a vertical orientation, and only 8.1% were round, with little variation across the subtypes. In mammogram, 35 (26.7%) cases were classified as benign,

mostly due to circumscribed mass without calcification (11.5%) (Table II and Figure 3). Table II shows the univariate associations between imaging features and TNBC subtypes. In ultrasound, BL and IM subtypes were more often spiculated (61.2% and 68.4%) than ML and LAR (46.7% and 37.8%). However, the differences were not found to be statistically significant. Around 80% of tumors exhibited hypoechoic pattern. A possible relationship between sonographic posterior pattern and subtype was observed, with less posterior enhancement in LAR (29.7% versus 56.7-57.9%).

BL-TNBC presented more often as a mass without calcification in mammogram than other subtypes (71.4% versus 48.6-57.9%). Mammographic margins of LAR-TNBC were more often spiculated (24.3% versus 0-4.1% in other subtypes).




Figure 1. The relationship of tumor shape in ultrasound to initial BI-RADS classification (benign/probably benign versus probably malignant).




Figure 2. Ultrasound images of BL-TNBC with lobulated margins (A) and IM-TNBC with spiculated margins (B).

The relationship of ultrasound and mammogram findings to initial BI-RADS classification (benign/probably benign *versus* probably malignant) is displayed in Figures 1 and 4. In few cases, BI-RADS classification 1 to 3 was given despite “typical” malignant patterns such as mass with calcification in mammogram or irregular shape in ultrasound.

## Discussion

Due to the molecular heterogeneity of TNBC, a marker-expression based subclassification has been established over the last few years. Several recent studies have detected different oncogenic alterations and underline the fact that




Figure 3. Mammogram of ML-TNBC with round shape and circumscribed margin.

TNBC can be further stratified which may lead to a targeted therapy in TNBC (11). Currently, standard treatment for TNBC includes chemotherapy. In many cases, neoadjuvant chemotherapy has the advantages of observing the responsiveness in-vivo, minimizing the surgical procedure, and providing an early start to the systemic therapy. As drug responsiveness differs within the TNBC subtypes, early subclassification in the neoadjuvant setting can be helpful in therapeutic decisions (12, 13). Not only pathological characteristics from a diagnostic biopsy specimen, but also characteristics such as growth pattern in clinical imaging of TNBC subtypes will play an important role in identifying the most promising therapy for every patient.

TNBC is known to present a benign appearance more often in mammogram and ultrasound than non-TNBC (7, 14). Due to the heterogeneous pattern of TNBC especially in ultrasound, some authors have concluded that the predictive value of sonographic appearance might be low when comparing TNBC *versus* non-TNBC (7). However, the histological subtype heterogeneity of TNBC that may explain a broad variety of clinical imaging characters has rarely been considered so far (15, 16). In our study, we compared four histological subtypes of TNBC: BL, IM, ML, and LAR. This subtype classification is more and more used in the current literature; however, data about clinical relevance is still poor (3, 5, 17). Our study is the first that compares the radiological features of four TNBC subtypes.

We observed more often suspicious signs such as spiculated margins or a mass with calcifications in mammogram of LAR subtype. These results are in line with recent studies (18, 19). In these studies, histologic grade did not differ between LAR-TNBC and control group, whereas we observed a lower percentage of poorly differentiated tumors in our group of LAR-TNBC. Lower histological grading and slower growth can be associated with a more spiculated, infiltrating growth pattern, giving the surrounding tissue time for stromal interactions (7). This may explain our mammographic findings; however, they seem to be contradictory to the more frequent lobular sonographic patterns of LAR in our study. In these cases, it is possible that the tumor was overseen in mammogram or showed indistinct margins in the correlating mammogram. Smooth, pushing borders in clinical imaging as well as in the correlating histopathological microscopic pictures are typical for highly aggressive tumors (Figure 5). However, our results imply that not only the grade of differentiation, but also the histopathological subtype of TNBC has an impact on its presentation in ultrasound and mammogram.

BI-RADS classification provides an established algorithm to distinguish benign and potential malignant lesions in the breast. However, daily practice shows a more complex interaction of clinical presentation, risk factors, correlation between mammogram and ultrasound, and individual examiner's pattern recognition. This may explain some initial misclassifications by the first examiner of patients in our study group despite retrospective review of the images occasionally revealing suspicious signs as shown in Figures 1 and 4.

Our study has some limitations to discuss. It is a retrospective single centre study, and despite the reviewers of the clinical images being blinded for the histopathological TNBC subtype, they knew the diagnosis of malignancy. On the other hand, a review process has the advantage of providing a more objective evaluation of imaging features. Furthermore, it allows an interpretation of image characteristics leading to potential misclassification of malignant tumors.

The heterogeneous group of TNBC also shows a great variety of clinical imaging appearance. The authors believe that the presented analysis, as well as further studies, will help to provide an early clinical-pathological subtyping of TNBC prior to a systemic therapy. This is even more relevant, as developing targeted therapies for TNBC subtypes is the topic of latest and ongoing clinical trials (20). For example, if a newly diagnosed TNBC showed spiculated sonographic margins and mass with calcification in the mammogram, histopathological subtyping could be indicated to identify LAR. LAR has a lower pathologic complete response (pCR) to chemotherapy, which might influence therapeutic decisions.




Figure 4. The relationship of tumor shape in mammogram to initial BI-RADS classification (benign/probably benign versus probably malignant).

Further studies with higher numbers of subtyped TNBC are warranted to elaborate the impact of histopathological subtype and tumor differentiation on radiological imaging.

### Conflicts of Interest

The Authors declare that they have no competing interests regarding this study.

### Authors' Contributions

All Authors made substantial contributions to the study, they have approved the current version and agreed on publication.

### Acknowledgements

The Authors are grateful to all patients who participated in this research.

### Funding

This research was self-funded by the Breast Center Zurich and the Institute of Pathology and Molecular Pathology, University Hospital of Zurich.




Figure 5. Microscopic slide (H&E stain) of a surgical specimen of BL-TNBC that presented benign characteristics in ultrasound. Arrows show the smooth, "pushing" border of the tumor.

## References

- Dent R, Trudeau M, Pritchard KI, Hanna WM, Kahn HK, Sawka CA, Lickley LA, Rawlinson E, Sun P and Narod SA: Triple-negative breast cancer: clinical features and patterns of recurrence. *Clin Cancer Res* 13: 4429-4434, 2007. PMID: 17671126. DOI: 10.1158/1078-0432.CCR-06-3045
- Kaplan HG and Malmgren JA: Impact of triple negative phenotype on breast cancer prognosis. *Breast J* 14: 456-463, 2008. PMID: 18657139. DOI: 10.1111/j.1524-4741.2008.00622.x
- Lehmann BD, Bauer JA, Chen X, Sanders ME, Chakravarthy AB, Shyr Y and Pietenpol JA: Identification of human triple-negative breast cancer subtypes and preclinical models for selection of targeted therapies. *J Clin Invest* 121: 2750-2767, 2011. PMID: 21633166. DOI: 10.1172/JCI45014.
- Le Du F, Eckhardt BL, Lim B, Litton JK, Moulder S, Meric-Bernstam F, Gonzalez-Angulo AM and Ueno NT: Is the future of personalized therapy in triple-negative breast cancer based on molecular subtype? *Oncotarget* 6: 12890-12908, 2015. PMID: 25973541. DOI: 10.18632/oncotarget.3849
- Turner NC and Reis-Filho JS: Tackling the diversity of triple-negative breast cancer. *Clin Cancer Res* 19: 6380-6388, 2013. PMID: 24298068. DOI: 10.1158/1078-0432.CCR-13-0915.
- Balleuquier C, Ayadi S, van Nguyen K, Vanel D, Dromain C and Sigal R: BIRADS classification in mammography. *Eur J Radiol* 61: 192-194, 2007. PMID: 17164080. DOI: 10.1016/j.ejrad.2006.08.033
- Li JW, Zhang K, Shi ZT, Zhang X, Xie J, Liu JY and Chang C: Triple-negative invasive breast carcinoma: the association between the sonographic appearances with clinicopathological feature. *Sci Rep* 8: 9040, 2018. PMID: 29899425. DOI: 10.1038/s41598-018-27222-6
- Gagliato DM, Gonzalez-Angulo AM, Lei X, Theriault RL, Giordano SH, Valero V, Hortobagyi GN and Chavez-Macgregor M: Clinical impact of delaying initiation of adjuvant chemotherapy in patients with breast cancer. *J Clin Oncol* 32: 735-744, 2014. PMID: 24470007. DOI: 10.1200/JCO.2013.49.7693
- Kündig P, Giesen C, Jackson H, Bodenmiller B, Papassotiropoulos B, Freiburger SN, Aquino C, Opitz L and Varga Z: Limited utility of tissue micro-arrays in detecting intra-tumoral heterogeneity in stem cell characteristics and tumor progression markers in breast cancer. *Transl Med* 16: 118, 2018. PMID: 29739401. DOI: 10.1186/s12967-018-1495-6
- Elfgen C, Reeve K, Moskovsky L, Güth U, Bjelic-Radisic V, Fleisch M, Tausch CJ and Varga Z: Prognostic impact of PIK3CA protein expression in triple negative breast cancer and its subtypes. *J Cancer Res Clin Oncol* 145: 2051-2059, 2019. PMID: 31270600. DOI: 10.1007/s00432-019-02968-2
- Azim HA, Ghosn M, Oualla K and Kassem L: Personalized treatment in metastatic triple negative breast cancer: The outlook in 2020. *Breast J* 26: 69-80, 2019. PMID: 31872557. DOI: 10.1111/tbj.13713
- Hill DP, Harper A, Malcolm J, McAndrews MS, Mockus SM, Patterson SE, Reynolds T, Baker EJ, Bult CJ, Chesler EJ and Blake JA: Cisplatin-resistant triple-negative breast cancer subtypes: multiple mechanisms of resistance. *BMC Cancer* 19: 1039, 2019. PMID: 31684899. DOI: 10.1186/s12885-019-6278-9
- Lehmann BD, Jovanovic B, Chen X, Estrada MV, Johnson KN, Shyr Y, Moses HL, Sanders ME and Pietenpol JA: Refinement of triple-negative breast cancer molecular subtypes: Implications for neoadjuvant chemotherapy selection. *PLoS One* 11, 2016. PMID: 27310713. DOI: 10.1371/journal.pone.0157368
- Gao B, Zhang H, Zhang SD, Cheng XY, Zheng SM, Sun YH, Zhang DW, Jiang Y and Tian JW: Mammographic and clinicopathological features of triple-negative breast cancer. *Br J Radiol* 87: 1039, 2014. PMID: 24734934. DOI: 10.1259/bjr.20130496
- Boissierie-Lacroix M, Macgrogan G, Debled M, Ferron S, Asad-Syed M, McKelvie-Sebileau P, Mathoulin-Pelissier S, Brouste V and Hurtevent-Labrot G: Triple-negative breast cancers: associations between imaging and pathological findings for triple-negative tumors compared with hormone receptor-positive/human epidermal growth factor receptor-2-negative breast cancers. *Oncologist* 18: 802-811, 2013. PMID: 23821326. DOI: 10.1634/theoncologist.2013-0380
- Krizmanich-Conniff KM, Paramagul C, Patterson SK, Helvie MA, Roubidoux MA, Myles JD, Jiang K and Sabel M: Triple receptor-negative breast cancer: imaging and clinical characteristics. *AJR Am J Roentgenol* 199: 458-464, 2012. PMID: 22826413. DOI: 10.2214/AJR.10.6096
- Li Z, Ren M, Tian J, Jiang S, Liu Y, Zhang L, Wang Z, Song Q, Liu C and Wu T: The differences in ultrasound and clinicopathological features between basal-like and normal-like subtypes of triple negative breast cancer. *PLoS One* 10(3), 2015. PMID: 25734578. DOI: 10.1371/journal.pone.0114820
- Bae MS, Park SY, Song SE, Kim WH, Lee SH, Han W, Park IA, Noh DY and Moon WK: Heterogeneity of triple-negative breast cancer: mammographic, US, and MR imaging features according to androgen receptor expression. *Eur Radiol* 25: 419-427, 2015. PMID: 25224727. DOI: 10.1007/s00330-014-3419-z
- Candelaria RP, Adrada BE, Wei W, Thompson AM, Santiago L, Lane DL, Huang ML, Arribas EM, Rauch GM, Symans WF, Gilcrease MZ, Huo L, Lim B, Ueno NT, Moulder SL and Yang WT: Imaging features of triple-negative breast cancers according to androgen receptor status. *Eur J Radiol* 114: 167-174, 2019. PMID: 31005169. DOI: 10.1016/j.ejrad.2019.03.017
- Damaskos C, Garmpi A, Nikolettos K, Vavourakis M, Diamantis E, Patsouras A, Farmaki P, Nonni A, Dimitroulis D, Mantas D, Antoniou EA, Nikolettos N, Kontzoglou K and Garmpis N: Triple-negative breast cancer: the progress of targeted therapies and future tendencies. *Anticancer Res* 39: 5285-5296, 2019. PMID: 31570423. DOI: 10.21873/anticancer.13722

Received February 23, 2020

Revised March 8, 2020

Accepted March 13, 2020